



## AMENDMENTS TO THE CLAIMS

Claim 1 (Currently Amended): A method for selectively inhibiting the C-terminal site of angiotensin I converting enzyme comprising utilizing administering to a patient in need thereof at least one phosphinic pseudopeptide derivative comprising the amino acid sequence of formula (I) below:



wherein,


-  $R_2$  and  $R_3$ , which are identical or different, represent the side chain of a natural or unnatural amino acid, the sequence:



also possibly forming the Pro (proline) residue, and


-  $R_5$  represents a hydrogen atom, a pharmacologically acceptable counterion, or a group that ~~can form~~ forms an *in vivo* hydrolysable phosphinic ester  $\frac{1}{2}$ .

Claim 2 (Currently Amended): A method for selectively inhibiting the C-terminal site of angiotensin I converting enzyme comprising ~~utilizing~~ administering to a patient in need thereof a phosphinic pseudopeptide derivative corresponding to formula (II) below:



wherein,

- R<sub>1</sub> represents a protecting group for an amine function, or an amino acid or a peptide protected with a protecting group for an amine function,
- R<sub>2</sub> and R<sub>3</sub>, which may be identical or different, represent the side chain of a natural or unnatural amino acid, the sequence:



also possibly forming the Pro residue,

- R<sub>4</sub> represents a hydrogen atom or a pharmacologically acceptable counterion,
- and
- R<sub>5</sub> represents a hydrogen atom, a pharmacologically acceptable counterion, or a group that can form forms an *in vivo* hydrolysable phosphinic ester ; .

Claim 3 (Previously Presented): The method of Claim 2, wherein  $R_1$  represents a protecting group for an amine function chosen from acetyl and benzyloxycarbonyl groups.

Claim 4 (Previously Presented): The method of Claim 1, wherein  $R_2$  represents the benzyl, methyl or phenylethyl group.


Claim 5 (Previously Presented): The method of Claim 1, wherein  $R_3$  represents the side chain of alanine, arginine or tryptophan.

Claim 6 (Currently Amended): The method of Claim 1, wherein the sequence  $-\text{NH}-\text{CH}(\text{R}_3)-\text{CO}-$  forms the Pro residue:




Claim 7 (Previously Presented): The method of Claim 1, wherein  $R_4$  and/or  $R_5$  represent(s) a hydrogen atom.

Claim 8 (Currently Amended): The method of Claim 2, wherein the phosphinic  
pseudopeptide derivative corresponds to the formula is:




(pseudo-peptide G)

Claim 9 (Currently Amended): A phosphinic pseudopeptide derivative comprising the amino acid sequence of formula (I) below:



wherein,

- R<sub>2</sub> represents the side chain of a natural or unnatural amino acid,
- the sequence:




forms the Pro residue:



- ~~R<sub>4</sub> represents a hydrogen atom or a pharmacologically acceptable counterion,~~
- and
- R<sub>5</sub> represents a hydrogen atom, a pharmacologically acceptable counterion, or a group ~~that can form~~ that forms an *in vivo* hydrolysable phosphinic ester.

Claim 10 (Currently Amended): A phosphinic pseudopeptide derivative  
corresponding to of formula (II) below:




wherein,

- R<sub>1</sub> represents a protecting group for an amine function, or an amino acid or a peptide protected with a protecting group for an amine function,

- R<sub>2</sub> represents the side chain of a natural or unnatural amino acid,

- the sequence:



forms the Pro residue:




- R<sub>4</sub> represents a hydrogen atom or a pharmacologically acceptable counterion,

and

- R<sub>5</sub> represents a hydrogen atom, a pharmacologically acceptable counterion, or a group ~~that can form~~ that forms an *in vivo* hydrolysable phosphinic ester.


Claim 11 (Currently Amended): A phosphinic pseudopeptide derivative of formula:



(pseudo-peptide G)


Claim 12 (Previously Presented): A pharmaceutical composition comprising at least one phosphinic pseudopeptide derivative as claimed in Claim 9.

Claim 13 (Currently Amended): A pharmaceutical composition, ~~in which the~~ comprising a phosphinic pseudopeptide derivative corresponds to the of formula:




(pseudo-peptide G)

Claim 14 (Currently Amended): A process for preparing a pseudopeptide of formula:



wherein:

- $R_1$  represents a protecting group for an amine function, or an amino acid or a peptide protected with a protecting group for an amine function,
- $R_2$  and  $R_3$ , which may be identical or different, represent the side chain of a natural or unnatural amino acid, the sequence:




also possibly forming the Pro residue, and

- $R_4$  and  $R_5$  represent a hydrogen atom;


which comprises the following steps:

- 1) reacting a compound of formula (III):




in which  $R_1$  and  $R_2$  are as defined above, with the compound of formula (IV):







in which Ac represents the acetyl group and Et represents the ethyl group, to obtain the compound of formula (V):





2) converting compound (V) into compound (VI) by reacting compound (V) with sodium borohydride:



3) protecting the hydroxyl group of compound (VI) with a protecting group  $R^5$ , for example the adamantyl group Ad, to give the compound of formula (VII):





4) saponifying compound (VII) to give the compound of formula (VIII):



5) coupling the compound of formula (VIII) with the amino acid of formula (IX)

or (X):




in which  $\text{R}_3$  is as defined above, and

6) removing the protecting group  $\text{Ad}$  and  $\text{R}^5$ .


Claim 15 (Previously Presented): A process as claimed in Claim 14, wherein the peptide coupling step 5) is performed via solid-phase peptide synthesis wherein the solid phase is a resin substituted with the amino acid of formula (IX) or (X).

Claim 16 (Currently Amended): A process for preparing a pseudopeptide of formula:



wherein,

- R<sub>1</sub> represents a protecting group for an amine function, or an amino acid or a peptide protected with a protecting group for an amine function,
- R<sub>2</sub> and R<sub>3</sub>, which may be identical or different, represent the side chain of a natural or unnatural amino acid, the sequence:




also possibly forming the Pro residue,

- R<sub>4</sub> represents a hydrogen atom, and
- R<sub>5</sub> represents a group that ~~can form~~ forms an *in vivo* hydrolysable phosphinic ester;

wherein the phosphinic function of the pseudopeptide obtained via the process of Claim 14 is esterified by coupling with an alcohol of formula R<sub>5</sub>OH or by reaction with a halide of formula R<sub>5</sub>X in which X represents a halogen atom.

Claim 17 (Currently Amended): A compound of formula (VIII):



wherein:

- Ad represents an adamantyl group.

-  $R_1$  represents a protecting group for an amine function or an amino acid or a peptide protected with an amine function, and

-  $R_2$  represents the side chain of a natural or unnatural amino acid.

Claim 18 (Previously Presented): The method of Claim 2, wherein  $R_2$  represents the benzyl, methyl or phenylethyl group.

Claim 19 (Previously Presented): The method of Claim 2, wherein  $R_3$  represents the side chain of alanine, arginine or tryptophan.

Claim 20 (Currently Amended): The method of Claim 2, wherein the sequence  $-\text{NH}-\text{CH}(\text{R}_3)-\text{CO}-$  forms the Pro residue:




Claim 21 (Previously Presented): The method of Claim 2, wherein  $R_4$  and/or  $R_5$  represent(s) a hydrogen atom.

Claim 22 (Previously Presented): A pharmaceutical composition comprising at least one phosphinic pseudopeptide derivative as claimed in Claim 10.


Claim 23 (Previously Presented): A pharmaceutical composition comprising at least one phosphinic pseudopeptide derivative as claimed in Claim 11.

Claim 24 (Currently Amended): A process for preparing a pseudopeptide of formula:



wherein,

- R<sub>1</sub> represents a protecting group for an amine function, or an amino acid or a peptide protected with a protecting group for an amine function,
- R<sub>2</sub> and R<sub>3</sub>, which may be identical or different, represent the side chain of a natural or unnatural amino acid, the sequence:



also possibly forming the Pro residue,

- R<sub>4</sub> represents a hydrogen atom, and
- R<sub>5</sub> represents a group that can form forms an *in vivo* hydrolysable phosphinic ester;

wherein the phosphinic function of the pseudopeptide obtained via the process of Claim 15 is esterified by coupling with an alcohol of formula R<sub>5</sub>OH or by reaction with a halide of formula R<sub>5</sub>X in which X represents a halogen atom.

Claim 25 (New): A process as claimed in Claim 14, wherein R<sup>5</sup> is an adamantyl group.